



Federal Reserve Bank of Chicago

**Simpler Bootstrap Estimation of the
Asymptotic Variance of U-statistic
Based Estimators**

Bo E. Honoré and Luojia Hu

August 2015

WP 2015-07

Simpler Bootstrap Estimation of the Asymptotic Variance of U-statistic Based Estimators*

Bo E. Honoré[†] Luojia Hu[‡]

August 4, 2015

*This research was supported by the National Science Foundation and the Gregory C. Chow Econometric Research Program at Princeton University and presented at the Federal Reserve Bank of Chicago. The opinions expressed here are those of the authors and not necessarily those of the Federal Reserve Bank of Chicago or the Federal Reserve System. We are thankful to Reka Zempleni for excellent research assistance. The most recent version of this paper can be found at <http://www.princeton.edu/~honore/papers/RiccatiBootstrap.pdf>.

[†]Department of Economics, Princeton University, Princeton, NJ 08544-1021. Email: honore@princeton.edu.

[‡]Economic Research Department, Federal Reserve Bank of Chicago, 230 S. La Salle Street, Chicago, IL 60604. Email: llhu@frbchi.org.

Abstract

The bootstrap is a popular and useful tool for estimating the asymptotic variance of complicated estimators. Ironically, the fact that the estimators are complicated can make the standard bootstrap computationally burdensome because it requires repeated re-calculation of the estimator. In Honoré and Hu (2015) we propose a computationally simpler bootstrap procedure based on repeated re-calculation of one-dimensional estimators. The applicability of that approach is quite general. In this paper, we propose an alternative method which is specific to extremum estimators based on U -statistics. The contribution here is that rather than repeated re-calculating the U -statistic-based estimator, we can recalculate a related estimator based on single-sums. A simulation study suggests that the approach leads to a good approximation to the standard bootstrap, and that if this is the goal, then our approach is superior to numerical derivative methods.

KEYWORDS: U -statistics; bootstrap; inference; numerical derivatives.

JEL CODE: C10, C18.

1 Introduction

Applied researchers often use the bootstrap when estimating standard errors. There are a number of advantages of using the bootstrap for estimating standard errors. One is that it is computationally convenient because it requires little additional programming above and beyond what is required for calculating the estimator. A second is that direct estimation of the asymptotic variance of estimators might require choosing tuning parameters such as bandwidths for nonparametric estimation or step-lengths for numerical derivatives. See Hong, Mahajan, and Nekipelov (2015) for a discussion of this. On the other hand, the bootstrap can be computationally demanding because it requires one to recalculate the same estimator many times. This is especially true for estimators that are based on double-sums such as those in Honoré and Powell (1994) and Alan, Honoré, Hu, and Leth-Petersen (2014).

The contribution of this paper is to point out that for extremum estimators based on U -statistics, one can sometimes estimate the Hessian term of the asymptotic variance by a simpler bootstrap procedure which involves repeated re-estimation of an extremum estimator based on simple averages. This will lower the computational burden of the bootstrap without giving up the aforementioned advantages of the bootstrap.

The proposed approach applies directly to classical rank-estimators of regression models such as those in Adichie (1967) and Sievers (1978) and to estimators of truncated and censored regression models like those in Bhattacharya, Chernoff, and Yang (1983), Honoré and Powell (1994) and Alan, Honoré, Hu, and Leth-Petersen (2014). The approach can also be used to estimate the Hessian-term in the asymptotic variance of rank correlation estimators like those in Han (1987), Cavanagh and Sherman (1998), Abrevaya (1999), Abrevaya (2000),

Khan and Tamer (2007) and Bhattacharya (2008).

Our approach relies heavily on the asymptotic variance of the estimator having the “sandwich”-structure, $H^{-1}VH^{-1}$, where both H and V are symmetric and positive definite. Interestingly, the approach does not apply to general U -statistic method-of-moments estimators because their asymptotic distribution will have a “sandwich”-structure without the restriction that H is symmetric and positive definite.

A small Monte Carlo study suggests that inference based on the proposed method will be more reliable than inference based on simple numerical derivatives, and that it performs as well as inference based on an infeasible second-order numerical derivative that uses the (unknown) optimal stepsize.

2 Background and Insight

Honoré and Powell (1994) derive the asymptotic properties for estimators $\hat{\theta}$ which minimize an m^{th} -order U-statistic of the form

$$U_n(\theta) = \binom{n}{m}^{-1} \sum_c p(z_{i_1}, z_{i_2}, \dots, z_{i_m}, \theta) \quad (1)$$

where z_i is the data for observation i and the sum is taken over the $\binom{n}{m}$ combinations of m distinct elements $\{i_1, \dots, i_m\}$ from the set $\{1, \dots, n\}$. The “kernel” function $p(\cdot)$ is symmetric in its first m arguments and left- and right-differentiable in each component of θ in some open neighborhood of the true parameter θ_0 .

The estimator $\hat{\theta}$ satisfies an approximate first-order condition of the form

$$Q_n(\hat{\theta}) \equiv \sqrt{n} \binom{n}{m}^{-1} \sum_c q(z_{i_1}, \dots, z_{i_m}, \hat{\theta}) = o_p(1),$$

where each component of the vector $q(\cdot)$ is a convex combination of the left and right partial derivatives of p with respect to the corresponding component of θ . Assuming random sampling, the resulting asymptotic distribution of θ has a limiting normal distribution:

$$\sqrt{n}(\hat{\theta} - \theta_0) \xrightarrow{d} N(0, H_0^{-1} V_0 H_0^{-1}), \quad (2)$$

with

$$V_0 \equiv m^2 E[r(z_i, \theta_0) r(z_i, \theta_0)'] \quad \text{and} \quad H_0 \equiv \partial E[r(z_i, \theta_0)] / \partial \theta'$$

where $r(z_i, \theta) \equiv E[q(z_i, z_{i_2}, \dots, z_{i_m}, \theta) | z_i]$.

Consistent estimators of the matrices H_0 and V_0 are needed to carry out large-sample inference on θ_0 . Estimation of V_0 is unproblematic. Define an estimator of $r(z_i, \theta)$ by

$$\hat{r}(z_i, \theta) \equiv \binom{n-1}{m-1}^{-1} \sum_{c'} q(z_i, z_{i_2}, \dots, z_{i_m}, \theta),$$

where the sum is now taken over the $\binom{n-1}{m-1}$ combinations of $m-1$ distinct elements $\{i_2, \dots, i_m\}$ from the set $\{1, \dots, n\} \setminus \{i\}$. An estimator of V_0 which uses $\hat{r}(z_i, \hat{\theta})$ is

$$\hat{V}_n \equiv \frac{m^2}{n} \sum_{i=1}^n \hat{r}(z_i, \hat{\theta}) \hat{r}(z_i, \hat{\theta})'.$$

Estimation of H_0 is more problematic because the function $q(\cdot, \theta)$ is discontinuous in many interesting examples. When this is the case, the explicit expression for H_0 will involve densities. One possibility is to estimate these nonparametrically. Alternatively, as noted by Pakes and Pollard (1989), one can use a “numerical derivative” estimator of H_0 even when $q(\cdot, \theta)$ is not differentiable. Specifically, an estimator of the ℓ th column of H_0 is given by

$$\hat{H}_{n\ell} \equiv (2\hat{h})^{-1} \left(Q_n(\hat{\theta} + \hat{h}e_\ell) - Q_n(\hat{\theta} - \hat{h}e_\ell) \right), \quad (3)$$

where e_ℓ is the ℓ th standard basis vector and $\hat{h}$ is a (possibly stochastic) sequence of bandwidths satisfying $\hat{h} = o_p(1)$ and $(\sqrt{n}\hat{h})^{-1} = O_p(1)$. See also Hong, Mahajan, and Nekipelov (2015).

Although these estimators of H_0 will be consistent, they are somewhat unattractive because of their dependence on tuning parameters such as bandwidths for nonparametric regression and step-lengths for numerical derivatives. In cases like this, many researchers prefer estimating the variance via a bootstrap. Unfortunately, when m is greater than 1, this is often computationally infeasible because of the large number of terms in (1)¹. We therefore introduce an alternative approach to estimating H_0 .

The new insight exploited in this paper is that one can bootstrap the variance of an alternative estimator with the same H -matrix and then use that variance estimator to back out an estimator of the variance of the estimator of interest. Specifically, consider an alternative “split-sample” estimator, where the minimand involves an average of only $n^* = \text{int}(n/m)$ terms using the same kernel evaluated at “non-overlapping” m -tuples of observations; i.e., the minimand is of the form

$$U_n^*(\theta) \equiv \frac{1}{n^*} \sum_{i=1}^{n^*} p(z_i, z_{i+n^*}, \dots, z_{i+(m-1)n^*}, \theta). \quad (4)$$

Since the terms in this minimand are i.i.d. for each θ , the usual consistency and asymptotic normality arguments for M -estimators apply and we have

$$\sqrt{n}(\tilde{\theta} - \theta_0) \xrightarrow{d} N(0, H_1^{-1} \Omega H_1^{-1}), \quad (5)$$

¹Subsampling is also unlikely to be feasible in practice unless the subsamples are very small, in which case one would worry about whether the asymptotics provide good finite sample approximations.

with $\Omega \equiv mV \left[q \left(z_i, z_{i+n^*}, \dots, z_{i+(m-1)n^*}, \theta_0 \right) \right]$ and, importantly, $H_1 = H_0$ under random sampling. Note that it is again easy to estimate Ω .

Since minimization of (4) only involves n^* terms, it is typically quite simple to estimate the variance of $\tilde{\theta}$ via the bootstrap. Letting $Avar \left(\tilde{\theta} \right) = H_0^{-1} \Omega H_0^{-1}$, this suggests estimating H_0^{-1} by solving the equation

$$\widehat{Avar} \left(\tilde{\theta} \right) = X \hat{\Omega}_n X \quad (6)$$

for a symmetric and positive definite X .

Equations like (6) are special cases of what is known as the continuous time algebraic Riccati equation. See Lancaster and Rodman (1995). Ricatti equations play an important role in control theory, and results exist that guarantee uniqueness of positive definite solutions. It is also well understood how to solve Riccati equations numerically. See, for example, Ni (2008) and Arnold and Laub (1984).

The approach pursued here could also be used to estimate the ‘‘Hessian’’ for maximum rank correlation-like estimators like those in Han (1987), Cavanagh and Sherman (1998), Abrevaya (1999), Abrevaya (2000), Khan and Tamer (2007) and Bhattacharya (2008). For these U-statistics estimators, the naive split-sample estimators would be an estimator similar to the maximum score estimator of Manski (1975). However, this would not be directly fruitful because this estimator is not asymptotically normally distributed (Cavanagh (1987) and Kim and Pollard (1990)). On the other hand, smoothed versions of the maximum score estimator like the one in Horowitz (1992) will be asymptotically normal with the same Hessian as the maximum rank correlation estimator, although the rate of convergence will be different. The reason why we do not pursue this approach here, is that the matrices in

the middle of the variance expressions in (2) and (5) (V and Ω) depend on densities. So estimating them would require tuning parameters such as bandwidths or number of nearest neighbors.

More general just-identified method-of-moments estimators also have “sandwich” asymptotic variance of the form $H^{-1}VH'^{-1}$. The main difference is that in this case, H will typically not be symmetric and positive definite. We therefore can not rely on the results about uniqueness of solutions to Riccati equations. Indeed, it is not difficult to see that equations of the form $\widehat{Avar}(\tilde{\theta}) = X\widehat{\Omega}_nX'$ do not have unique solutions. For example, both $\begin{pmatrix} \sqrt{1+\sqrt{3/4}} & \frac{1}{2}(\sqrt{1+\sqrt{3/4}})^{-1} \\ \frac{1}{2}(\sqrt{1+\sqrt{3/4}})^{-1} & \sqrt{1+\sqrt{3/4}} \end{pmatrix}$ and $\begin{pmatrix} \sqrt{2} & 0 \\ \sqrt{1/2} & \sqrt{3/2} \end{pmatrix}$ will be solutions when $\widehat{Avar}(\tilde{\theta}) = \begin{pmatrix} 2 & 1 \\ 1 & 2 \end{pmatrix}$ and $\widehat{V}_1 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$. In other words, for method-of-moments estimators, it is not possible to identify the “Hessian”-term, H^{-1} , from the equation $\Sigma = H^{-1}VH'^{-1}$.

3 Monte Carlo Illustration

We illustrate the usefulness of the approach by conducting a small Monte Carlo study of one of the pairwise comparison estimators in Honoré and Powell (1994). In this section we will implicitly assume that the researcher considers the bootstraps the “gold standard” and that the goal is to compare it with more easily computable alternatives. Of course, in practice, the behavior of test-statistics also depends on how well the finite sample distribution of the estimator is approximated by its limiting normal distribution. It is possible that the estimation error of a poor variance estimator can offset this approximation error and lead to better finite sample performance of the resulting test-statistics. For this reason, we will think of the bootstrap as the benchmark in the discussion below.

Honoré and Powell (1994) propose estimators of censored and truncated regression models. Here we focus on the censored regression model defined by $y_i = \max(0, x_i'\beta + \varepsilon_i)$ where ε_i is independent of x_i . Honoré and Powell (1994) showed that subject to regularity conditions, β is the unique minimizer of $E[s(y_i, y_j, (x_i - x_j)'b)]$ where

$$s(y_1, y_2, \delta) = \begin{cases} \Xi(y_1) - (y_2 + \delta)\xi(y_1) & \text{for } \delta < -y_2 \\ \Xi(y_1 - y_2 - \delta) & \text{for } -y_2 \leq \delta \leq y_1 \\ \Xi(-y_2) - (\delta - y_1)\xi(-y_2) & \text{for } y_1 < \delta \end{cases}$$

and $\Xi'(d) = \xi(d)$ (for almost all d) for some convex function Ξ . Natural choices for Ξ are $\Xi(d) = d^2$ and $\Xi(d) = |d|$.

They therefore define an estimator of the censored regression model by minimization of

$$S_n(b) = \binom{n}{2}^{-1} \sum_{i < j} s(y_i, y_j, (x_i - x_j)'b). \quad (7)$$

Note that $S_n(b)$ is a convex function of b . $S_n(b)$ is differentiable provided that Ξ is. If Ξ is not differentiable at 0, then the points of non-differentiability of $S_n(b)$ occur when $y_i - y_j - (x_i - x_j)'b = 0$ and either $y_i \neq 0$ or $y_j \neq 0$ or both.

This section presents the results from a Monte Carlo experiment designed to investigate the performance of the variance estimator proposed here. We consider the estimator proposed in Honoré and Powell (1994) with $\Xi(d) = d^2$ and $\Xi(d) = |d|$. When $\Xi(d) = d^2$ it is easy to estimate the asymptotic variance of the estimator without having to choose a bandwidth. We include it to illustrate the performance of our approach in a setting where we expect it to perform relatively poorly compared to other methods.

There are four explanatory variables. There are all dependent and have the following

marginal distributions. x_1 is binary, x_2 is chi-square, x_3 is normal with probability $1/2$ and 0 otherwise, and x_4 is normal. All the variables are normalized to have mean 0 and variance 1 and the correlation structure for the four variables is constructed as follows. First we generate x^* from a four-dimensional normal distribution with means 0, variances 1 and correlations 0.5. We then define $x_1 = 2 \cdot 1(x_1^* > 0) - 1$; $x_2 = (x_2^{*2} - 1)/\sqrt{2}$, $x_3 = \sqrt{2}x_3^*D$ where D is independent of x^* and Bernoulli with $p = \frac{1}{2}$. Finally $x_4 = x_4^*$.

The error, ε_i , is independent of the other variables and t-distributed with three degrees of freedom and β is a vector of ones. This gives an R^2 in an uncensored version of the model of approximately 0.48. There is no additional constant in the generating process. This results in a censoring probability of approximately 51%.

The Monte Carlo experiment has 400 replications. All bootstraps are performed using 1000 bootstrap samples. In each replication, we calculate the estimators defined by minimizing (7) with $\Xi(d) = d^2$ and with $\Xi(d) = |d|$. When $\Xi(d) = d^2$, $S_n(b)$ is almost everywhere twice differentiable so there is no particular reason to use the approach proposed here because one can use analytical derivatives. We estimate the variance of $\hat{\beta}$ by the bootstrap, the Riccati bootstrap proposed here and by estimating H by (two-sided) numerical derivatives of the analytical derivative of $s(y_i, y_j, (x_i - x_j)'b)$ using stepsizes $h_\ell = 0.05/\text{std}(x_\ell)$, $h_\ell = 0.25/\text{std}(x_\ell)$ and $h_\ell = 0.5/\text{std}(x_\ell)$. The bootstrap draws for the Riccati bootstrap are independent of the draws for the regular bootstrap. The numerical derivative estimator of H is $\frac{1}{2}(\hat{H} + \hat{H}')$ where $\hat{H}$ is given in (3).² When $\Xi(d) = d^2$, we also estimate H by its

²We make the estimator of H symmetric to ensure that $\hat{H}^{-1}\hat{V}\hat{H}^{-1}$ is positive definite (provided that $\hat{H}$ is invertible). Alternatively we could have constructed $\hat{H}^{-1}\hat{V}(\hat{H}^{-1})'$ or $(\hat{H}^{-1})'\hat{V}\hat{H}^{-1}$.

sample analog. Since variance estimation is typically used for inference, we first calculate the “ t -test” statistic, T , for whether each coefficient equals its true value using each of the variance estimators. We then construct $\Phi(T)$ where Φ is the normal cumulative distribution function. This corresponds to the p -value for a one-sided test. Since the aim of this paper is to construct a more convenient alternative to the bootstrap variance, we compare the p -values using the different variance estimators with those obtained using the bootstrap variance.

It is well understood that asymptotic normality does not imply that moments are finite. We therefore use a robust estimator for the bootstrap variances. Specifically, with a sample of B bootstrap estimators, $\tilde{\beta}_1, \dots, \tilde{\beta}_B$, we estimate the variance of the ℓ 'th coordinate, $\tilde{\beta}(\ell)$, by the square of 1.34 times the interquartile range of $\tilde{\beta}_1(\ell), \dots, \tilde{\beta}_B(\ell)$. To estimate the correlation matrix, we transform the ℓ 'th coordinate of $\tilde{\beta}_b(\ell)$ into $\Phi^{-1}\left(\frac{R_B(\tilde{\beta}_b(\ell))}{B+1}\right)$ where $R_B(\tilde{\beta}_b(\ell))$ is the rank of $\tilde{\beta}_b(\ell)$. We then use the sample correlation of $\Phi^{-1}\left(\frac{R_B(\tilde{\beta}_b(\ell))}{B+1}\right)$ to estimate the correlation matrix of $\tilde{\beta}$. If the empirical distribution of $\tilde{\beta}$ is closely approximated by a normal distribution, then this correlation matrix will be very similar to the empirical correlation of $\tilde{\beta}$. The advantage is that the former is much less sensitive to outliers.

Tables 1 and 2 report the Monte Carlo average of the absolute difference in percentage points between the p -values based on any pair of the variance estimates averaged over all four parameters when $n = 100$. Table 1 corresponds to $\Xi(d) = d^2$ while Table 2 shows the results when $\Xi(d) = |d|$. In addition, they also include estimators of the variance based on non-robust bootstrap and Riccati bootstrap variance estimation.

The results in Table 1 are all expected. The objective function that defines the estimator

is almost everywhere twice differentiable, so there is almost no difference between calculating the Hessian and approximating it numerically. This is especially true when the stepsize is small. Both the bootstrap and Riccati bootstrap yield p -values that are within 2 to 3 percentage points of each other and of those from the analytical and numerical derivative estimators. The only outlier is the Riccati bootstrap estimator based on non-robust estimation of the variance. This is not surprising since the estimator based on $\Xi(d) = d^2$ is related to the OLS estimator in a linear regression model with fairly fat tails.³ Table 2 is more interesting because $\Xi(d) = |d|$ gives an estimator for which we cannot estimate the Hessian term in the variance by analytical differentiation. In this case, the Riccati bootstrap does a much better job approximating the bootstrap than the numerical derivative estimators do.

The role of the stepsize in numerical differentiation is akin to the role of the bandwidth in nonparametric kernel estimation. This raises two considerations. One is that the results can be sensitive to the choice of stepsize, and the other is that it is possible to reduce bias by using higher order numerical derivatives. See for example Hong, Mahajan, and Nekipelov (2015). To investigate these, we calculate the numerical derivative estimator of the Hessian as above using stepsizes $h_\ell = h / \text{std}(x_\ell)$ for a range of h . In addition, using the notation from section 2, we consider the second order numerical derivative of Q_n defined by $\frac{1}{2} \left(\widetilde{H}_n + \widetilde{H}_n' \right)$

³Without censoring, the U -statistic estimator equals the slope of the OLS estimator of y on x and a constant, while the split-sample estimator would be the OLS estimator of the difference in the y on the difference in the x (without a constant). Since the errors are distributed according to a t -distribution with 3 degrees of freedom, these OLS estimators will have poorly-behaved sample moments.

where the ℓ 'th columns of $\tilde{H}_n$ is

$$\tilde{H}_{n\ell} \equiv \hat{h}^{-1} \left(-\frac{1}{12} Q_n \left(\hat{\theta} + 2\hat{h}e_\ell \right) + \frac{2}{3} Q_n \left(\hat{\theta} + \hat{h}e_\ell \right) - \frac{2}{3} Q_n \left(\hat{\theta} - \hat{h}e_\ell \right) + \frac{1}{12} Q_n \left(\hat{\theta} - 2\hat{h}e_\ell \right) \right).$$

Like the higher-order kernel estimators, this construction of $\tilde{H}_n$ removes the first order term in the bias of $\hat{H}_n$.

We calculate the average absolute difference between the p -values calculated from the bootstrap and the numerical derivative estimators using $\hat{H}_n$ and $\tilde{H}_n$ with stepsizes $h_\ell = h / \text{std}(x_\ell)$ for a range of h . These are presented in Figures 1-4 for sample sizes 50, 100, 200 and 400. The horizontal (fatter) line represents the Monte Carlo average of the absolute difference between the p -values calculated from the bootstrap and the Riccati bootstrap, while the thinner solid and the dashed lines represent the Monte Carlo average of the absolute difference between the p -values calculated from the bootstrap and those using $\tilde{H}_n$ and $\hat{H}_n$, respectively. The figures suggest that the Riccati bootstrap provides a better approximation to the bootstrap than approaches based on numerical derivative approximations to the Hessian regardless of the choices of stepsize.

4 Conclusion

The bootstrap is a popular and useful tool for estimating the asymptotic variance of complicated estimators. Unfortunately the standard bootstrap can be computationally burdensome because it requires repeated re-calculation of the estimator. In Honoré and Hu (2015) we proposed a modification to the standard bootstrap that replaces re-estimation of the original parameter vector with re-estimation of one-dimensional estimators. In this paper, we

focus on estimators based on minimization of U -statistics. Our contribution here is to replace re-estimation of the original parameter vector with re-estimation of estimators based on minimization of simple averages. We perform a small Monte Carlo experiment for an estimator whose asymptotic variance involves unknown functions. In this case, the common alternative to the standard bootstrap is to estimate the Hessian in the asymptotic variance by numerical derivatives. We find that our approach performs better than this alternative regardless of the choice of tuning parameter of the numerical derivative.

References

- ABREVAYA, J. (1999): “Leapfrog Estimation of a Fixed-Effects Model with Unknown Transformation of the Dependent Variable,” *Journal of Econometrics*, 93, 203–228.
- (2000): “Rank Estimation of a Generalized Fixed Effects Regression Model,” *Journal of Econometrics*, 95, 1–23.
- ADICHIE, J. N. (1967): “Estimates of Regression Parameters Based on Rank Tests,” *The Annals of Mathematical Statistics*, 38(3), pp. 894–904.
- ALAN, S., B. E. HONORÉ, L. HU, AND S. LETH-PETERSEN (2014): “Estimation of Panel Data Regression Models with Two-Sided Censoring or Truncation,” *Journal of Econometric Methods*, 3(1), 1–20.
- ARNOLD, W.F., I., AND A. LAUB (1984): “Generalized eigenproblem algorithms and software for algebraic Riccati equations,” *Proceedings of the IEEE*, 72(12), 1746–1754.

- BHATTACHARYA, D. (2008): “A Permutation-Based Estimator for Monotone Index Models,” *Econometric Theory*, 24(3), pp. 795–807.
- BHATTACHARYA, P. K., H. CHERNOFF, AND S. S. YANG (1983): “Nonparametric Estimation of the Slope of a Truncated Regression,” *The Annals of Statistics*, 11(2), pp. 505–514.
- CAVANAGH, C. (1987): “The Limiting Behavior of Estimators Defined by Optimization,” Harvard University.
- CAVANAGH, C. L., AND R. P. SHERMAN (1998): “Rank Estimators of Monotonic Index Models,” *Journal of Econometrics*, 84, 351–381.
- HAN, A. (1987): “Nonparametric Analysis of a Generalized Regression Model,” *Journal of Econometrics*, 35, 303–316.
- HONG, H., A. MAHAJAN, AND D. NEKIPELOV (2015): “Extremum estimation and numerical derivatives,” *Journal of Econometrics*, 188(1), 250 – 263.
- HONORÉ, B. E., AND L. HU (2015): “Poor (Wo)man’s Bootstrap,” *FRB of Chicago Working Paper No. 2015-01*. Available at SSRN: <http://ssrn.com/abstract=2577508> or <http://dx.doi.org/10.2139/ssrn.2577508>.
- HONORÉ, B. E., AND J. L. POWELL (1994): “Pairwise Difference Estimators of Censored and Truncated Regression Models,” *Journal of Econometrics*, 64(1-2), 241–78.
- HOROWITZ, J. L. (1992): “A Smoothed Maximum Score Estimator for the Binary Response Model,” *Econometrica*, 60, 505–531.

- KHAN, S., AND E. TAMER (2007): “Partial rank estimation of duration models with general forms of censoring,” *Journal of Econometrics*, 136(1), 251 – 280.
- KIM, J., AND D. POLLARD (1990): “Cube Root Asymptotics,” *Ann. Statist.*, 18(1), 191–219.
- LANCASTER, P., AND L. RODMAN (1995): *Algebraic Riccati Equations*. Oxford University Press, USA.
- MANSKI, C. F. (1975): “The Maximum Score Estimation of the Stochastic Utility Model of Choice,” *Journal of Econometrics*, 3, 205–228.
- NI, M.-L. (2008): “Existence Condition on Solutions to the Algebraic Riccati Equation,” *Acta Automatica Sinica*, 34(1), 85 – 87.
- PAKES, A., AND D. POLLARD (1989): “Simulation and the Asymptotics of Optimization Estimators,” *Econometrica*, 57, 1027–1057.
- SIEVERS, G. L. (1978): “Weighted Rank Statistics for Simple Linear Regression,” *Journal of the American Statistical Association*, 73(363), pp. 628–631.

Table 1: Difference in p -values in Percentage Points Using Different Variance Estimators.

$$\Xi(d) = d^2, n = 100.$$

	Analyt.	Boots.	Riccati	$h = 0.05$	$h = 0.25$	$h = 0.50$	Boot(NR)	Ricc.(NR)
Analy.	0.00	1.25	2.77	0.07	0.17	0.36	2.17	6.28
Boots.	1.25	0.00	2.05	1.25	1.23	1.20	1.39	5.37
Riccati	2.77	2.05	0.00	2.77	2.76	2.73	1.79	3.83
$h = 0.05$	0.07	1.25	2.77	0.00	0.15	0.35	2.17	6.27
$h = 0.25$	0.17	1.23	2.76	0.15	0.00	0.23	2.15	6.26
$h = 0.50$	0.36	1.20	2.73	0.35	0.23	0.00	2.10	6.24
Boot(NR)	2.17	1.39	1.79	2.17	2.15	2.10	0.00	4.36
Ricc.(NR)	6.28	5.37	3.83	6.27	6.26	6.24	4.36	0.00

Table 2: Difference in p -values in Percentage Points Using Different Variance Estimators.

$$\Xi(d) = |d|, n = 100.$$

	Analyt.	Boots.	Riccati	$h = 0.05$	$h = 0.25$	$h = 0.50$	Boot(NR)	Ricc.(NR)
Analyt.	NA	NA	NA	NA	NA	NA	NA	NA
Boots.	NA	0.00	1.67	4.66	4.28	4.34	1.10	2.24
Riccati	NA	1.67	0.00	4.05	3.60	3.82	1.64	1.26
$h = 0.05$	NA	4.66	4.05	0.00	2.03	2.25	4.23	3.41
$h = 0.25$	NA	4.28	3.60	2.03	0.00	0.89	3.78	2.78
$h = 0.50$	NA	4.34	3.82	2.25	0.89	0.00	3.72	2.81
Boot(NR)	NA	1.10	1.64	4.23	3.78	3.72	0.00	1.63
Ricc.(NR)	NA	2.24	1.26	3.41	2.78	2.81	1.63	0.00

Figure 1: Relative Performance of Numerical Derivatives vs. Riccati Bootstrap. $n = 50$




Figure 2: Relative Performance of Numerical Derivatives vs. Riccati Bootstrap. $n = 100$




Figure 3: Relative Performance of Numerical Derivatives vs. Riccati Bootstrap. $n = 200$





Figure 4: Relative Performance of Numerical Derivatives vs. Riccati Bootstrap. $n = 400$



Working Paper Series

A series of research studies on regional economic issues relating to the Seventh Federal Reserve District, and on financial and economic topics.

Examining Macroeconomic Models through the Lens of Asset Pricing <i>Jaroslav Borovička and Lars Peter Hansen</i>	WP-12-01
The Chicago Fed DSGE Model <i>Scott A. Brave, Jeffrey R. Campbell, Jonas D.M. Fisher, and Alejandro Justiniano</i>	WP-12-02
Macroeconomic Effects of Federal Reserve Forward Guidance <i>Jeffrey R. Campbell, Charles L. Evans, Jonas D.M. Fisher, and Alejandro Justiniano</i>	WP-12-03
Modeling Credit Contagion via the Updating of Fragile Beliefs <i>Luca Benzoni, Pierre Collin-Dufresne, Robert S. Goldstein, and Jean Helwege</i>	WP-12-04
Signaling Effects of Monetary Policy <i>Leonardo Melosi</i>	WP-12-05
Empirical Research on Sovereign Debt and Default <i>Michael Tomz and Mark L. J. Wright</i>	WP-12-06
Credit Risk and Disaster Risk <i>François Gourio</i>	WP-12-07
From the Horse's Mouth: How do Investor Expectations of Risk and Return Vary with Economic Conditions? <i>Gene Amromin and Steven A. Sharpe</i>	WP-12-08
Using Vehicle Taxes To Reduce Carbon Dioxide Emissions Rates of New Passenger Vehicles: Evidence from France, Germany, and Sweden <i>Thomas Klier and Joshua Linn</i>	WP-12-09
Spending Responses to State Sales Tax Holidays <i>Sumit Agarwal and Leslie McGranahan</i>	WP-12-10
Micro Data and Macro Technology <i>Ezra Oberfield and Devesh Raval</i>	WP-12-11
The Effect of Disability Insurance Receipt on Labor Supply: A Dynamic Analysis <i>Eric French and Jae Song</i>	WP-12-12
Medicaid Insurance in Old Age <i>Mariacristina De Nardi, Eric French, and John Bailey Jones</i>	WP-12-13
Fetal Origins and Parental Responses <i>Douglas Almond and Bhashkar Mazumder</i>	WP-12-14

Working Paper Series *(continued)*

Repos, Fire Sales, and Bankruptcy Policy <i>Gaetano Antinolfi, Francesca Carapella, Charles Kahn, Antoine Martin, David Mills, and Ed Nosal</i>	WP-12-15
Speculative Runs on Interest Rate Pegs The Frictionless Case <i>Marco Bassetto and Christopher Phelan</i>	WP-12-16
Institutions, the Cost of Capital, and Long-Run Economic Growth: Evidence from the 19th Century Capital Market <i>Ron Alquist and Ben Chabot</i>	WP-12-17
Emerging Economies, Trade Policy, and Macroeconomic Shocks <i>Chad P. Bown and Meredith A. Crowley</i>	WP-12-18
The Urban Density Premium across Establishments <i>R. Jason Faberman and Matthew Freedman</i>	WP-13-01
Why Do Borrowers Make Mortgage Refinancing Mistakes? <i>Sumit Agarwal, Richard J. Rosen, and Vincent Yao</i>	WP-13-02
Bank Panics, Government Guarantees, and the Long-Run Size of the Financial Sector: Evidence from Free-Banking America <i>Benjamin Chabot and Charles C. Moul</i>	WP-13-03
Fiscal Consequences of Paying Interest on Reserves <i>Marco Bassetto and Todd Messer</i>	WP-13-04
Properties of the Vacancy Statistic in the Discrete Circle Covering Problem <i>Gadi Barlevy and H. N. Nagaraja</i>	WP-13-05
Credit Crunches and Credit Allocation in a Model of Entrepreneurship <i>Marco Bassetto, Marco Cagetti, and Mariacristina De Nardi</i>	WP-13-06
Financial Incentives and Educational Investment: The Impact of Performance-Based Scholarships on Student Time Use <i>Lisa Barrow and Cecilia Elena Rouse</i>	WP-13-07
The Global Welfare Impact of China: Trade Integration and Technological Change <i>Julian di Giovanni, Andrei A. Levchenko, and Jing Zhang</i>	WP-13-08
Structural Change in an Open Economy <i>Timothy Uy, Kei-Mu Yi, and Jing Zhang</i>	WP-13-09
The Global Labor Market Impact of Emerging Giants: a Quantitative Assessment <i>Andrei A. Levchenko and Jing Zhang</i>	WP-13-10

Working Paper Series *(continued)*

Size-Dependent Regulations, Firm Size Distribution, and Reallocation <i>François Gourio and Nicolas Roys</i>	WP-13-11
Modeling the Evolution of Expectations and Uncertainty in General Equilibrium <i>Francesco Bianchi and Leonardo Melosi</i>	WP-13-12
Rushing into the American Dream? House Prices, the Timing of Homeownership, and the Adjustment of Consumer Credit <i>Sumit Agarwal, Luojia Hu, and Xing Huang</i>	WP-13-13
The Earned Income Tax Credit and Food Consumption Patterns <i>Leslie McGranahan and Diane W. Schanzenbach</i>	WP-13-14
Agglomeration in the European automobile supplier industry <i>Thomas Klier and Dan McMillen</i>	WP-13-15
Human Capital and Long-Run Labor Income Risk <i>Luca Benzoni and Olena Chyruk</i>	WP-13-16
The Effects of the Saving and Banking Glut on the U.S. Economy <i>Alejandro Justiniano, Giorgio E. Primiceri, and Andrea Tambalotti</i>	WP-13-17
A Portfolio-Balance Approach to the Nominal Term Structure <i>Thomas B. King</i>	WP-13-18
Gross Migration, Housing and Urban Population Dynamics <i>Morris A. Davis, Jonas D.M. Fisher, and Marcelo Veracierto</i>	WP-13-19
Very Simple Markov-Perfect Industry Dynamics <i>Jaap H. Abbring, Jeffrey R. Campbell, Jan Tilly, and Nan Yang</i>	WP-13-20
Bubbles and Leverage: A Simple and Unified Approach <i>Robert Barsky and Theodore Bogusz</i>	WP-13-21
The scarcity value of Treasury collateral: Repo market effects of security-specific supply and demand factors <i>Stefania D'Amico, Roger Fan, and Yuriy Kitsul</i>	WP-13-22
Gambling for Dollars: Strategic Hedge Fund Manager Investment <i>Dan Bernhardt and Ed Nosal</i>	WP-13-23
Cash-in-the-Market Pricing in a Model with Money and Over-the-Counter Financial Markets <i>Fabrizio Mattesini and Ed Nosal</i>	WP-13-24
An Interview with Neil Wallace <i>David Altig and Ed Nosal</i>	WP-13-25

Working Paper Series *(continued)*

Firm Dynamics and the Minimum Wage: A Putty-Clay Approach <i>Daniel Aaronson, Eric French, and Isaac Sorkin</i>	WP-13-26
Policy Intervention in Debt Renegotiation: Evidence from the Home Affordable Modification Program <i>Sumit Agarwal, Gene Amromin, Itzhak Ben-David, Souphala Chomsisengphet, Tomasz Piskorski, and Amit Seru</i>	WP-13-27
The Effects of the Massachusetts Health Reform on Financial Distress <i>Bhashkar Mazumder and Sarah Miller</i>	WP-14-01
Can Intangible Capital Explain Cyclical Movements in the Labor Wedge? <i>François Gourio and Leena Rudanko</i>	WP-14-02
Early Public Banks <i>William Roberds and François R. Velde</i>	WP-14-03
Mandatory Disclosure and Financial Contagion <i>Fernando Alvarez and Gadi Barlevy</i>	WP-14-04
The Stock of External Sovereign Debt: Can We Take the Data at ‘Face Value’? <i>Daniel A. Dias, Christine Richmond, and Mark L. J. Wright</i>	WP-14-05
Interpreting the <i>Pari Passu</i> Clause in Sovereign Bond Contracts: It’s All Hebrew (and Aramaic) to Me <i>Mark L. J. Wright</i>	WP-14-06
AIG in Hindsight <i>Robert McDonald and Anna Paulson</i>	WP-14-07
On the Structural Interpretation of the Smets-Wouters “Risk Premium” Shock <i>Jonas D.M. Fisher</i>	WP-14-08
Human Capital Risk, Contract Enforcement, and the Macroeconomy <i>Tom Krebs, Moritz Kuhn, and Mark L. J. Wright</i>	WP-14-09
Adverse Selection, Risk Sharing and Business Cycles <i>Marcelo Veracierto</i>	WP-14-10
Core and ‘Crust’: Consumer Prices and the Term Structure of Interest Rates <i>Andrea Ajello, Luca Benzoni, and Olena Chyruk</i>	WP-14-11
The Evolution of Comparative Advantage: Measurement and Implications <i>Andrei A. Levchenko and Jing Zhang</i>	WP-14-12

Working Paper Series *(continued)*

Saving Europe?: The Unpleasant Arithmetic of Fiscal Austerity in Integrated Economies <i>Enrique G. Mendoza, Linda L. Tesar, and Jing Zhang</i>	WP-14-13
Liquidity Traps and Monetary Policy: Managing a Credit Crunch <i>Francisco Buera and Juan Pablo Nicolini</i>	WP-14-14
Quantitative Easing in Joseph's Egypt with Keynesian Producers <i>Jeffrey R. Campbell</i>	WP-14-15
Constrained Discretion and Central Bank Transparency <i>Francesco Bianchi and Leonardo Melosi</i>	WP-14-16
Escaping the Great Recession <i>Francesco Bianchi and Leonardo Melosi</i>	WP-14-17
More on Middlemen: Equilibrium Entry and Efficiency in Intermediated Markets <i>Ed Nosal, Yuet-Yee Wong, and Randall Wright</i>	WP-14-18
Preventing Bank Runs <i>David Andolfatto, Ed Nosal, and Bruno Sultanum</i>	WP-14-19
The Impact of Chicago's Small High School Initiative <i>Lisa Barrow, Diane Whitmore Schanzenbach, and Amy Claessens</i>	WP-14-20
Credit Supply and the Housing Boom <i>Alejandro Justiniano, Giorgio E. Primiceri, and Andrea Tambalotti</i>	WP-14-21
The Effect of Vehicle Fuel Economy Standards on Technology Adoption <i>Thomas Klier and Joshua Linn</i>	WP-14-22
What Drives Bank Funding Spreads? <i>Thomas B. King and Kurt F. Lewis</i>	WP-14-23
Inflation Uncertainty and Disagreement in Bond Risk Premia <i>Stefania D'Amico and Athanasios Orphanides</i>	WP-14-24
Access to Refinancing and Mortgage Interest Rates: HARPing on the Importance of Competition <i>Gene Amromin and Caitlin Kearns</i>	WP-14-25
Private Takings <i>Alessandro Marchesiani and Ed Nosal</i>	WP-14-26
Momentum Trading, Return Chasing, and Predictable Crashes <i>Benjamin Chabot, Eric Ghysels, and Ravi Jagannathan</i>	WP-14-27
Early Life Environment and Racial Inequality in Education and Earnings in the United States <i>Kenneth Y. Chay, Jonathan Guryan, and Bhashkar Mazumder</i>	WP-14-28

Working Paper Series *(continued)*

Poor (Wo)man's Bootstrap <i>Bo E. Honoré and Luojia Hu</i>	WP-15-01
Revisiting the Role of Home Production in Life-Cycle Labor Supply <i>R. Jason Faberman</i>	WP-15-02
Risk Management for Monetary Policy Near the Zero Lower Bound <i>Charles Evans, Jonas Fisher, François Gourio, and Spencer Krane</i>	WP-15-03
Estimating the Intergenerational Elasticity and Rank Association in the US: Overcoming the Current Limitations of Tax Data <i>Bhashkar Mazumder</i>	WP-15-04
External and Public Debt Crises <i>Cristina Arellano, Andrew Atkeson, and Mark Wright</i>	WP-15-05
The Value and Risk of Human Capital <i>Luca Benzoni and Olena Chyruk</i>	WP-15-06
Simpler Bootstrap Estimation of the Asymptotic Variance of U-statistic Based Estimators <i>Bo E. Honoré and Luojia Hu</i>	WP-15-07